

Buccolam[®] pre-filled syringes (Buccal Midazolam, 5mg/ml)

Parent and Carer's Information Leaflet and
Care Plan for Children at Risk of Long Seizures (Fits)



Individual Emergency Care Plan for:

Name.....DOB.....NHS no.....

Description of typical seizure when Buccolam[®] needs to be given:

.....

When should Buccolam[®] be given?.....

.....

Dose: Give:mg of Buccolam[®]. This is all of acoloured Buccolam[®] syringe.

Whenever you get a new prescription check that the concentration is 5mg/ml and the product is called Buccolam[®]. If it is not please contact us for a new care plan.

Additional Instructions?:.....

.....

When to call ambulance?.....

.....

Professional's Name..... **Signature**..... **Date**

I consent for other trained professionals to administer buccal midazolam as per this care plan.

Parent's Name.....**Signature**..... **Date**

For further advice contact your local team:

CEWT website

www.cewt.org.uk

Epilepsy Action Website:

www.epilepsy.org.uk

Young Epilepsy

www.youngepilepsy.org.uk

Many seizures will stop on their own. Some seizures need help to stop and may cause problems if they continue too long.

When do I give Buccolam®?

As well as this leaflet you should have had training on when and how to give Buccolam®. Buccolam® is the product name of a particular type of buccal Midazolam for your child. Generally Buccolam® may be given when a child or young person with epilepsy has:

- a generalised convulsive seizure lasting more than 5 minutes. (*These are seizures where the child does not respond, continues to be stiff and may jerk their arms and legs*). **OR**
- a pattern of convulsive seizures that stop then start again straight away without recovering in between and together go on longer than 5 minutes.
- Some children will have buccal midazolam for individual circumstances and this will be outlined on page 1 of this care plan if appropriate.

How do I store Buccolam®?

Buccolam® does not need to be stored in the fridge. Further details about storage can be found in the instruction leaflet supplied with the medication. Keep it out of reach of children. Check the expiry date before using. Ensure you take Buccolam® with you if you are going out. School or other carers may also need to keep some and arrange training.

How often can I use it?

When not in hospital, do not use Buccolam® more than once in any 24 hour period. Some children may have a plan describes more than once. see individual 'additional instructions'. If the first dose does not work, the ambulance or hospital may give a second dose 10 minutes after the first.

How do I measure the correct dose?



1. The buccal Midazolam will come in a box containing 4 tubes each with a pre-filled syringe.
2. There are different doses available so check you have the correct product and dose.

Red = 10mg;
Purple = 7.5mg;
Blue = 5mg;
Yellow = 2.5mg.

3. Remove the pre-filled syringe from



4. Check that the translucent tip-cap is also removed.



Translucent tip-cap

How do I give the Buccolam®?

1. If possible place your child on their side.
2. Place the tip of the syringe between the cheek and gums on the side closest to the floor.
3. Once in place, push the plunger down gently to give the medicine.
4. If your child is on their back give half the dose slowly in one side and then half in the other.



Does it work straight away?

Buccolam® will take 5 to 10 minutes to work and most children will sleep for a while after the seizure. They should be placed on their side during this time.

Do I need to call '999'?

- Generally, if your child has been given the buccal midazolam and is still having a seizure 10 minutes after giving the dose, **OR** if your child appears to have any difficulty breathing or stops breathing, call 999.
- If the child is needing buccal midazolam for the very first time, some carers may wish to first call 999 and then wait until the ambulance arrives before giving buccal midazolam
- Other parents/carers may wish to call an ambulance straight after giving the dose.

Are there any side effects?

Buccal Midazolam can make your child sleepy or restless. Very rarely it may slow down breathing.

Is it safe to give with other medicines?

Yes. But do not give it with rectal diazepam or other types of buccal Midazolam.

Test dose

Professionals may consider a test dose of buccal midazolam for some children where they have not had previous benzodiazepines and there are risk factors for an unexpected reaction.

Disclaimer

Some children have their own individual emergency plan which may differ from this leaflet. If you have any questions, please ask your doctor or specialist nurse.